WISCONSIN BEHAVIORAL RISK FACTOR SURVEY, 2002

The 2002 Wisconsin Behavioral Risk Factor Survey (BRFS) provides information based on responses from 4,356 adults (age 18 and older). Results are weighted to represent the 2.9 million adults ages 18-54, and 1.2 million age 55 and older, living in Wisconsin households.

PERCEIVED HEALTH STATUS

	All Adults		18-54		5	i5 +
	<u>%</u>	<u>(+/-)</u>	<u>%</u>	<u>(+/-)</u>	<u>%</u>	<u>(+/-)</u>
Health in general is:						
Excellent or very good	57	1	62	2	45	3
Good	31	1	30	2	34	
Fair or poor	12	1	8	1	21	2 2
Number of days during the						
past 30 that you felt						
sad, blue or depressed:						
0 days	65	1	62	2	71	2
1-5 days	25	1	27	2	19	
6-30 days	11	1	11	1	10	2 2
very healthy and full of						
energy:						
0 days	9	1	7	1	14	2
1-5 days	10	1	10	1	9	1
6-30 days	82	1	83	1	78	2
During past 30 days, for						
how many days did						
pain make it hard for						
you to do your usual						
activities:						
0 days	78	1	79	2	74	2
1-5 days	10	1	11	1	8	1
6-30 days	12	1	9	1	18	2

* See Technical Notes	*	See	Technical	Notes
-----------------------	---	-----	-----------	-------

TOBACCO	USE	OPIN	ION			
	All Adults		18-54		5	5 +
	<u>%</u>	<u>(+/-)</u>	<u>%</u>	<u>(+/-)</u>	<u>%</u>	<u>(+/-)</u>
Tobacco Use						
Current cigarette smoker *	23	1	28	2	13	2
Percent of smokers who quit smoking 1 day or longer in past year *	57	3	59	3	47	7
Former cigarette smoker	27	1	21	1	43	3

Tobacco Opinions *

Smoking in indoor work areas should be allowed:
...without restrictions
1 <1 1 1 1
...only in designated areas
29 2 30 2 25
...not at all
70 2 68 2 74

Smoking in restaurants should be allowed: ...without restrictions 1 <1 1 1 2 ...only in designated areas 47 2 50 3 41 ...not at all 51 2 49 3 57

OTHER RISK FACTORS FOR CHRONIC DISEASE

Overweight (BMI) # *	58	2	56	2	63	3
No leisure-time physical activity in past month	20	1	18	1	25	2
Servings of fruits and vegetables consumed per day:Less than 11-23-45 or more	5 36 36 23	1 1 1 1	7 40 34 20	1 2 2 1	2 25 41 32	1 2 3 2
Health professional has ever diagnosed:						
Asthma	12	1	12	1	11	2
Diabetes	5	1	3	1	11	2

** Excluding women who have had a hysterectomy

t Body Mass Index

FINDINGS FOR SELECTED YEARS							
	2002		1997		1992		
	<u>%</u>	(+/-)	<u>%</u>	(+/-)	<u>%</u>	<u>(</u> +/.	
Health Screenings							
In the past year had							
Pap smear (F)**	62	2	59	3	61	3	
clinical breast exam (F)	69	2	65	3	65	3	
mammogram (F 50+)	66	3	54	4	50	6	
Risk Behaviors							
Current cigarette smoker	23	1	23	2	23	2	
Males	25	2	26	3	23	3	
Females	21	2	21	2	23	3	
Overweight (BMI) ŧ *	58	2	55	2	52	3	
Males	68	2	65	3	63	4	
Females	48	2	46	3	41	3	
Always or nearly always use	82	1	81	2	78	2	
seatbelts when in a car Males	76	2	74	3	75	3	
Females	88	1	87	2	81	3	
During the past 30 days had 5 or more drinks on one occasion	25	1	23	2	25	2	
Males	36	2	36	3	34	3	
Females	14	1	11	2	17	3	
During the past 30 days drove after having too much to drink	4	1	5	1	5	1	
Males	7	1	9	2	7	2	
Females	2	1	2	1	3	1	

^{(+/-) = 95%} confidence interval

TECHNICAL NOTES

The Behavioral Risk Factor Survey is a representative, statewide telephone survey of Wisconsin household residents 18 years and older. Results are based on self-reports from respondents who are selected through a stratified random sample. Results are weighted to account for both nonresponse and sample design, and to adjust for the age-sex distribution of Wisconsin's population. The survey was designed by the Bureau of Health Information in cooperation with the U.S. Centers for Disease Control and Prevention. Survey sampling and interviewing were conducted by the University of Wisconsin Survey Center.

In 2002, 2,866 respondents were 18 to 54 years old, and 1,463 were 55 or older. There were 2,245 respondents in 1997 and 1,528 in 1992.

Columns labeled (+/-) indicate the 95% confidence interval associated with each estimate. For example, for all adults whose health is excellent or very good (57%), there is a 95% chance that the actual percentage falls within the range of plus or minus 1 percentage point (56%-58%).

The Centers for Disease Control and Prevention provided the following definitions:

- Overweight is defined by the Body Mass Index (BMI), a calculation (weight in kilograms divided by height in meters squared) using selfreported height and weight. A BMI greater than 25.0 is considered overweight.
- Current smokers are those who smoke currently, either every day or only some days, and have smoked more than 100 cigarettes in their lifetime.
- The percentage of **smokers who quit for 1 day or longer in past year** is based on those who smoke every day (those who smoke only some days are excluded).
- A drink of alcohol is defined as one can or bottle of beer, one glass of wine, one can or bottle of wine cooler, one cocktail, or one shot of liquor.
- The prevalence of **diabetes** is likely to be an underestimate because a significant proportion of adults have never been tested.
- Tobacco opinion questions were asked only from July to December.

Differences in estimates between years, age groups, or sexes may not be statistically significant. For additional information about the BRFSS, call Anne Ziege at 608-267-9821. For other health data, visit our Web site at http://dhfs.wisconsin.gov/stats/index.htm. Copies of this report are available at http://dhfs.wisconsin.gov/stats/BRFS.htm.

Health Counts in Wisconsin

New Findings from the Bureau of Health Information

BEHAVIORAL RISK FACTORS

2002

Division of Health Care Financing Department of Health and Family Services